

EXTENDED LY α EMISSION AROUND YOUNG QUASARS: A CONSTRAINT ON GALAXY FORMATION

ZOLTÁN HAIMAN¹

Princeton University Observatory, Princeton, NJ 08544, USA
 zoltan@astro.princeton.edu

MARTIN J. REES

Institute of Astronomy, Madingley Road, Cambridge, CB3 0HA, UK
 mjr@ast.cam.ac.uk

Draft version February 1, 2008

ABSTRACT

The early stage in the formation of a galaxy inevitably involves a spatially extended distribution of infalling, cold gas. If a central luminous quasar turned on during this phase, it would result in significant extended Ly α emission, possibly accompanied by other lines. For halos condensing at redshifts $3 \lesssim z \lesssim 8$ and having virial temperatures $2 \times 10^5 \text{ K} \lesssim T_{\text{vir}} \lesssim 2 \times 10^6 \text{ K}$, this emission results in a “fuzz” of characteristic angular diameter of a few arcseconds, and surface brightness $\sim 10^{-18} - 10^{-16} \text{ erg s}^{-1} \text{ cm}^{-2} \text{ asec}^{-2}$. The fuzz around bright, high redshift quasars could be detected in deep narrow band imaging with current telescopes, providing a direct constraint on galaxy formation models. The absence of detectable fuzz might suggest that most of the protogalaxy’s gas settles to a self-gravitating disk before a quasar turns on. However, continued gas infall from large radii, or an on-going merger spreading cold gas over a large solid angle, during the luminous quasar phase could also result in extended Ly α emission, and can be constrained by deep narrow band imaging.

Subject headings: cosmology: theory – galaxies: formation – quasars: general – black hole physics

1. INTRODUCTION

There has been considerable progress recently in the study of the formation of galaxies and quasars. Both the galaxy (Steidel et al. 1999) and quasar (Fan et al. 2000a) luminosity functions are now observationally determined to redshifts of $z \sim 4$, probing into the epoch when at least some of these objects are still expected to be assembling. Although the evolutions of the galaxy and quasar populations is generally expected to be connected, the detailed nature of this link is yet to be elucidated. One possibility is that quasars represent a brief phase in the early life of each galaxy. Indeed, theoretically, one might expect quasar activity to be triggered by the mergers that a galaxy is experiencing during its assembly (e.g. Cavaliere & Vittorini 2000, Kauffmann & Haehnelt 2000). This assertion has some observational support: the galaxy abundance appears to peak at a somewhat later redshift than does the quasar abundance (e.g. Madau & Pozzetti 2000); and at least infrared-selected quasars appear to preferentially reside in morphologically disturbed hosts (e.g. Baker & Clements 1997).

It is therefore interesting to consider a quasar that turns on within an assembling protogalaxy. The behavior of the gas inside a dark matter (DM) condensation during the formation of disk galaxies has been investigated in semi-analytic schemes (White & Rees 1978; Fall & Efstathiou 1980; Mo, Mao & White 1998; van den Bosch & Dalcanton 2000), and numerical simulations (Katz 1991; Navarro & White 1994; Navarro, Frenk & White 1997 [NFW]; Navarro & Steinmetz 1997; Moore et al. 1999). In these studies, the bulk of the baryons in the halo cool and settle into a rotationally supported disk. In the simplest picture,

the disk material originates as smooth gas, collapsing from the virial radius R_{vir} to its final orbital radius of $\sim \lambda R_{\text{vir}}$, where $\lambda \sim 0.05$ is the typical spin parameter. Numerical simulations have revealed a more complex process, where a fraction of the infalling gas forms smaller clumps early on; these clumps then progressively merge together, collide, and dissipate to form larger systems.

A robust feature of galaxy formation is, at least in the early stages, a spatially extended distribution of gas. At the densities expected at $z > 2$ for this gas ($\gtrsim 100$ times the background density), the radiative cooling time-scales are shorter than the typical dynamical times. In the absence of heat input from stars or a quasar, a significant fraction of this gas would therefore be cold ($T \lesssim 10^4 \text{ K}$) and neutral (Fall & Rees 1985). As it contracts inside the DM halo, the cold gas is heated by the halo potential; this heat is dissipated largely via collisional excitation of the Ly α , and possible other (metal) lines. The resulting line radiation results in a potentially observable, extended, low-surface brightness Ly α “fuzz” (Haiman et al. 2000; Fardal et al. 2000).

The presence of a bright central quasar during this phase could strongly enhance the surface brightness in the Ly α line, since a significant fraction of the quasar’s ionizing radiation could be reprocessed into recombination radiation in the same line. The purpose of this paper is to quantify the expected Ly α fluxes, and describe constraints that the presence or absence of extended Ly α “fuzz” around luminous quasars implies for the host galaxy. This paper is organized as follows: in § 2, we describe a simple toy-model for the distribution of cold gas in a protogalaxy; in § 3, we characterize the resulting Ly α “fuzz”; in § 4, we

¹Hubble Fellow

discuss the implied constraints on galaxy formation; and in § 5, we summarize our conclusions. Throughout this work, we adopt a Λ CDM cosmology with $\Omega_m = 0.3$, $\Omega_b = 0.04$, $\Omega_\Lambda = 0.7$, $h = 0.7$, and $\sigma_8 = 0.9$.

2. THE AMOUNT OF COLD GAS: A SPHERICAL MODEL

In this section, we describe a simple model for the structure of a spherically symmetric, two-phase gas in a DM halo without a central ionizing source. We shall assume that the gas has a centrally condensed radial profile $\rho(r)$, and that it consists of a two-phase medium with a cold ($T \sim 10^4$ K) mass fraction $f(r)$, and a hot ($T \sim T_{\text{vir}}$) mass fraction $1 - f(r)$. The physical state of the gas in a realistic protogalaxy is, of course, likely to be much more complicated, with asymmetric infall of pre-existing dense clumps, and the settling of gas in the central regions into a disk. Nevertheless, this model will serve as a reference point for our discussion in § 3 and § 4 below.

For sake of concreteness, we assume that the gas initially settles to a radial profile $\rho_{\text{gas}}(r)$ that satisfies hydrostatic equilibrium within a dark matter halo. For the profile of the halo, we follow the description in Navarro, Frenk & White (1997) with a concentration parameter $c = 5$. However, we assume that the average enclosed mass density is a fraction Δ_c of the critical density, where we obtain $\Delta_c \lesssim 18\pi^2$ from the spherical top-hat collapse (rather than using the fixed value of 200). To obtain $\rho_{\text{gas}}(r)$, we also assume that the gas is isothermal at the virial temperature of the halo (Makino, Sasaki & Suto 1998). It is useful to note that under these assumptions, the gas is centrally condensed with a flat core, and has a mean internal (volume-averaged) “clumping” of $\langle \rho_{\text{gas}}^2 \rangle \approx 2.8 \times 10^6 \rho_b^2$ relative to the background baryon density ρ_b .

In order to obtain the density $\rho_{\text{cold}}(r)$ of neutral gas, we next assume that a mass fraction $f(r)$ of the gas cools and condenses out into a cold phase with $T = 10^4$ K. Pressure balance with the remaining hot, ionized gas implies that the densities of the two components are

$$\eta_{\text{cold}} = \frac{\rho_{\text{cold}}}{\rho_{\text{gas}}} = f + (1 - f) \frac{T_{\text{vir}}}{10^4 \text{ K}} \geq 1$$

$$\eta_{\text{hot}} = \frac{\rho_{\text{hot}}}{\rho_{\text{gas}}} = (1 - f) + f \frac{10^4 \text{ K}}{T_{\text{vir}}} \leq 1. \quad (1)$$

The value of $f = f(r)$ is determined from the condition that the cooling time of the rarefied hot component is equal to the age of the system. The cooling time is given by $t_{\text{cool}} = (3/2)\mu m_p k_B T_{\text{vir}} / (\rho_{\text{hot}} \Lambda)$, where $\Lambda \sim 10^{-23} \text{ erg s}^{-1} \text{ cm}^3$ is the cooling function at T_{vir} . We conservatively adopt a metal-free cooling function (Böhringer & Hensler 1989). For the age of the system, we adopt 20% of the Hubble time, $t_{\text{age}} = 0.2(6\pi G \rho_z)^{-1/2}$, where ρ_z is the total (dark matter + baryons) mean background density at redshift z . This is roughly the “mass-doubling” time for halos of interest in the extended Press-Schechter formalism (see Lacey & Cole 1993; and Haiman, Spaans & Quataert 2000 for a discussion).

Illustrative results for the cold fraction $f(r)$ as a function of radius under the above assumptions are shown in Figure 1. The dashed and solid curves describe halos at redshifts $z = 3$, and $z = 5$, respectively. At both redshifts, four different halo sizes are shown, with virial temperatures of $T_{\text{vir}} = 2 \times 10^5 \text{ K}$, $4 \times 10^5 \text{ K}$, 10^6 K , and 10^7 K (top

to bottom). At $z = 3$, these correspond to halo masses $M_{\text{halo}} \approx 4 \times 10^{10} M_\odot$, $10^{11} M_\odot$, $5 \times 10^{11} M_\odot$, and $10^{13} M_\odot$; at $z = 5$, the halo masses are a factor of $\sim$ two smaller.

As expected, the cold fraction is a function of radius in each case, increasing towards $r = 0$ where densities are higher and cooling times are shorter. The cold fraction is larger for smaller halos, which have smaller initial binding energies, and hence cool more rapidly. The cold fraction increases with redshift, because of the higher densities and shorter cooling times. Although our model is highly idealized, it captures the above scalings (which are expected to be robust, as long as the cold fraction is determined by the cooling time), and provides a conservative estimate of the amount of cold gas. For comparison, we note that Mo & Miralda-Escudé (1996) have used a different, simplified model of a two-phase medium in order to model quasar line absorption systems. In their model, nearly all of the gas is cold within the “cooling radius”, defined as the radius at which the cooling time equals the Hubble time (exceeding R_{vir} for all halos considered here).




FIG. 1.— The mass fraction f of cold gas as a function of radius for two-phase gas inside a DM halo with an NFW profile. The dashed curves correspond to halos at redshift $z = 3$, and the solid curves to $z = 5$. At both redshifts, four different halo sizes are shown, corresponding (top to bottom) to, $T_{\text{vir}} = 2 \times 10^5 \text{ K}$, $4 \times 10^5 \text{ K}$, 10^6 K , and 10^7 K .

Finally, we emphasize that our model assumptions apply only within the virial radius, where the gas has been shock-heated. In the rest of this paper, we focus on the reprocessing of quasar light within protogalactic halos. However, the clumpy background gas outside the virial region can itself reprocess ionizing UV radiation into Ly α emission. Indeed, Gould & Weinberg (1996) have shown that Ly α absorption systems, illuminated by the UV background, can cause significant Ly α fluorescence (see Bunker et al. 1998 for a current status of observations). Numerical simulations suggest that the background gas near a collapsing protogalaxy is highly clumped, and hence could reprocess any ionizing radiation from the protogalaxy into Ly α emission. This type of reprocessing of quasar light could be the interpretation of extended Ly α –

emitting blobs found in the vicinity of quasars (Hu et al. 1991; Hu et al. 1996). Although these blobs are likely associated with small satellite protogalaxies, their Ly α emission could be dominated by reprocessed quasar light.

3. IONIZATION BY A CENTRAL SOURCE AND THE ACCOMPANYING LY α FUZZ

We next consider a quasar that turns on at the center of the halo described above. In general, the cold gas will then be photoionized by the quasar's UV radiation, out to a radius that depends on the quasar's ionizing luminosity. Utilizing the spatial distribution of cold gas obtained in our models, we first argue that reasonably-sized quasar black holes (BHs) can keep most of the cold phase photoionized. We then compute the characteristic Ly α surface brightness of such quasar-illuminated protogalaxies.

3.1. Effect of an Ionizing Source

The radius of the photoionized Strömgren sphere around a central quasar embedded in a spherical halo is given by

$$R_{\text{HII}} = \left[\frac{3\dot{N}_{\text{ph}}}{4\pi\alpha_B \langle n_{\text{H}}^2 \rangle} \right]^{1/3}, \quad (2)$$

where $\dot{N}_{\text{ph}}$ is the ionizing photon production rate of the central source, α_B is the hydrogen recombination coefficient evaluated at $\approx 10^4\text{K}$, and $\langle n_{\text{H}}^2 \rangle$ is the volume averaged mean squared density of cold hydrogen within R_{HII} . Note that in our models, the cold gas is compressed by a factor of η_{cold} , which enhances the local recombination rate by η_{cold}^2 . However, the cold gas occupies only a fraction f/η_{cold} of the volume, and hence the compression increases the total recombination rate by an overall factor of $f\eta_{\text{cold}}$.

Assuming a fixed cold mass fraction $f = 0.5$ across the halo, and assuming further that the central BH shines at the Eddington luminosity (for typical quasar spectra, this corresponds to an ionizing photon production rate of $\dot{N}_{\text{ph}} \approx 6 \times 10^{47}$ photons s^{-1} per $M_{\odot}$ of BH mass; see Cen & Haiman 2000), we then find the required size of the BH so that the Strömgren sphere extends all the way out to the virial radius:

$$M_{\text{bh}} \approx 6 \times 10^8 M_{\odot} \left(\frac{M_{\text{halo}}}{10^{12} M_{\odot}} \right)^{5/3} \left(\frac{1+z}{6} \right)^4. \quad (3)$$

Here we have utilized a relation between the halo mass, radius, and virial temperature from NFW. Equation (3) can be understood by recalling the scalings $R_{\text{HII}}^3 \propto M_{\text{bh}} \langle n_{\text{H}}^2 \rangle^{-1} \propto M_{\text{bh}} T_{\text{vir}}^{-1} (1+z)^{-6}$; $T_{\text{vir}} \propto M_{\text{halo}}^{2/3} (1+z)$; and $R_{\text{vir}} \propto M_{\text{halo}}^{1/3} (1+z)^{-1}$.

Equation (3) reveals that in most halos of interest ($3 \lesssim z \lesssim 8$; $10^{10} M_{\odot} \lesssim M_{\text{halo}} \lesssim 10^{12} M_{\odot}$), converting $\lesssim 2\%$ of the gas mass into a central BH is sufficient to keep most of the cold gas photoionized. This conclusion is conservative, since we assumed a constant cold fraction $f = 0.5$, which maximizes the internal clumping and the total recombination rate (cf. Eqn.1). Using the profiles of $f(r)$ obtained in our models, we have verified that for all halos considered in this work, the cold gas can be fully ionized by still smaller BHs, in all cases with $M_{\text{bh}} \leq 0.01 \times (\Omega_b/\Omega_m) M_{\text{halo}}$. Although the masses of BH's

in protogalaxies at $z \gtrsim 3$ are unknown, ratios as large as 1% of the gas mass would be consistent with the sizes of supermassive BH's found in nearby galaxies (Magorrian et al. 1998; Gebhardt et al. 2000; Ferrarese & Merritt 2000), especially since this ratio can evolve (and decrease) as a function of redshift (Menou, Haiman & Narayanan 2000).

3.2. Surface Brightness and Angular Size of Ly α Fuzz

Assuming that all of the cold gas in the model halo is photoionized, we obtain the total Ly α line luminosity as

$$L_{\alpha} = 0.68 \times E_{\alpha} \int_0^{R_{\text{vir}}} 4\pi r^2 dr \left(\frac{f}{\eta_{\text{cold}}} \right) n_{\text{H}}^2 \alpha_B, \quad (4)$$

where $E_{\alpha} = 10.2\text{eV}$ is the energy of a Ly α photon, and the integral represents the total recombination rate of photoionized gas within the halo. We have explicitly included the volume-filling factor (f/η_{cold}) of the cold gas, whose hydrogen number density is $n_{\text{H}} = 0.76\rho_{\text{cold}}/m_p$, and the fraction 0.68 of case B recombinations that yield a Ly α photon (Osterbrock 1989). We also note that additional recombination lines might be observable, including those of heavy elements, provided the halo gas is sufficiently pre-enriched (see De Breuck et al. 2000 for a review of relevant observations).




FIG. 2.— The characteristic surface brightness of Ly α fuzz around halos as a function of virial temperature. The solid curves correspond to redshifts $z = 3, 4, 5, 6, 7, 8$ (top to bottom near 10^7K). The dashed and dotted curves describe models (both at $z = 5$) in which the cooling time is increased by a factor of 10, or the age of the system is increased by a factor of 5, respectively.

For the halos shown in Figure 1, the values of L_{α} are $7 \times 10^{41} \text{ erg s}^{-1}$, $3 \times 10^{42} \text{ erg s}^{-1}$, $4 \times 10^{43} \text{ erg s}^{-1}$, and $3 \times 10^{46} \text{ erg s}^{-1}$ (at $z=3$), and $10^{42} \text{ erg s}^{-1}$, $5 \times 10^{42} \text{ erg s}^{-1}$, $5 \times 10^{43} \text{ erg s}^{-1}$, and $4 \times 10^{46} \text{ erg s}^{-1}$ (at $z=5$). The spatial extent of the Ly α -emitting gas is a fraction of the virial radius $R_{\text{vir}} \sim 10 - 100 \text{ kpc}$. As Figure 1 shows, this fraction varies inversely with the virial temperature of the halo. Here we define $R_{1/2}$ as the radius at which the cold fraction is $f = 0.5$, and assume that the Ly α radiation is emitted from within a characteristic radius, which

we take to be $R_\alpha \equiv \min(R_{1/2}, R_{\text{vir}})$. The characteristic angular size θ_α of the Ly α -emitting fuzz should then be $\sim \theta_\alpha = R_\alpha/d_A$, where d_A is the angular diameter distance. We find that for the halos shown in Figure 1, $\theta_\alpha \approx 2 - 3$ arcseconds. The angular size varies relatively little with redshift and virial temperature, and implies that the typical Ly α -emitting fuzz should appear extended and can be resolved with optical instruments.

The characteristic Ly α surface brightness within θ_α , $F_\alpha = L_\alpha/4\pi d_L^2/\pi\theta_\alpha^2$ (where d_L is the luminosity distance), is shown in Figure 2 as a function of T_{vir} . The solid curves correspond to redshifts $z = 3, 4, 5, 6, 7$, and 8 (top to bottom near 10^7K). This figure reveals several interesting features. First, the surface brightness is generally high, and all but the smallest halos should be potentially detectable with current instruments (see discussion below, and in Haiman, Spaans & Quataert 2000). Second, nearly all of the halo gas remains cold upto a virial temperature of $\sim 3 \times 10^5\text{K}$ (cf. Fig. 1). In this case, the Ly α fuzz extends out to the virial radius, and the surface brightness has the simple scaling $F_\alpha \propto \rho^2 V/R_{\text{vir}}^2(1+z)^4 \propto T_{\text{vir}}^{1/2}(1+z)^{1/2}$ (here $V \propto R_{\text{vir}}^3$ is the volume of the halo); i.e. the surface brightness weakly increases with both virial temperature and redshift. The latter result arises primarily from the strong dependence of the recombination rate on redshift, $\rho^2 \propto (1+z)^6$ [vs. the $(1+z)^4$ surface brightness dimming]. Third, for higher virial temperatures ($\gtrsim 3 \times 10^5\text{K}$), the cold fraction f falls somewhat below unity, allowing the cold phase to be compressed, and enhancing the recombination rate and the surface brightness. In this range of virial temperatures, the surface brightness scales approximately as $F_\alpha \propto T_{\text{vir}}^{5/2}(1+z)^{-2}$ [note that $\Lambda(T) \propto T^{1/2}$]. Both the stronger dependence on T_{vir} and the inverse scaling on redshift result from the dependence of the cold fraction on T_{vir} . Note that for still higher virial temperatures, the cold fraction would decrease to negligibly small values, and the surface brightness would drop sharply, but this happens only for exceedingly large halos.

In order to illustrate the robustness of the above conclusions, we have computed the surface brightness in two variants of our model. First, to facilitate comparison with earlier work, we adopt $t_{\text{cool}} = t_{\text{Hub}}$ rather than $t_{\text{cool}} = 0.2t_{\text{Hub}}$, as the condition used to compute the cold fraction f . The effect of this change on the surface brightness at $z = 5$ is shown by the dotted curve in Figure 2. As expected, the cold fraction remains unity for halos upto a higher virial temperature, which reduces the surface brightness by an order of magnitude at $T_{\text{vir}} \gtrsim 10^6\text{K}$. Second, we note that our simplified model would conflict with observations if we applied it to local galaxy clusters (see, e.g. Fabian 1994 for a review). Our model would predict a non-negligible amount of cold gas at the center of at least the lowest mass clusters (e.g. $M \sim 10^{14} M_\odot$ at redshift $z \sim 0$), where observations show little evidence for cold gas. This could mean that, in clusters, the cool gas converts quickly and efficiently into (low-mass) stars. Alternatively, there could be heat input (e.g. from higher-mass star formation, SNe, etc.) that effectively slows down the cooling. To mimic this latter scenario, we have increased the cooling time by a factor of 10, equivalent to requiring $t_{\text{cool}} = 0.02t_{\text{Hub}}$. This condition ensures that

$f \sim 0$ is predicted for all $z \sim 0$ galaxy clusters. The effect of this change (at $z = 5$) is shown by the dashed curve in Figure 2. The cold fraction is decreased, which enhances the surface brightness by upto an order of magnitude for halo with $T_{\text{vir}} \gtrsim 10^5\text{K}$.

4. OBSERVATIONAL PROSPECTS: CONSTRAINTS ON GALAXY AND QUASAR FORMATION

The main results of the previous two sections are (1) in galaxy-sized halos ($T_{\text{vir}} \gtrsim 10^6\text{K}$) at redshifts $z \gtrsim 3$, a significant fraction of the gas should be cold; and (2) if this gas is illuminated by ionizing radiation from a central quasar, it should be kept photoionized, and result in a detectable Ly α fuzz of characteristic size of $\gtrsim 2$ arcseconds, and surface brightness $\gtrsim 10^{-17} \text{ erg s}^{-1} \text{ cm}^{-2} \text{ asec}^{-2}$.

A handful of existing observations in various other contexts have already reached these relevant levels for the Ly α surface brightness. Examples are the imaging of a proto-cluster region of Lyman Break Galaxies at redshift $z \approx 3$ in a narrow-band Ly α filter, reaching a sensitivity of $\sim 10^{-17} \text{ erg s}^{-1} \text{ cm}^{-2} \text{ asec}^{-2}$ in a ~ 16 hour observation with the Palomar 200-inch telescope (Steidel et al. 2000). Similar sensitivities have been reached in ~ 5 hours with the Keck telescope in blank field searches for high-redshift Ly α galaxies (Cowie & Hu 1998), and should also be achievable by Subaru.² Extended Ly α emission has also been detected around high-redshift radio galaxies (e.g. De Breuck et al. 2000) and radio-loud quasars (possibly related to outflows; e.g. Heckman et al. 1991a,b; Bremer et al. 1992). Nevertheless, observations have not yet targeted bright quasars to search for extended Ly α emission at similar depths. Less sensitive observations (reaching a few $\times 10^{-16} \text{ erg s}^{-1} \text{ cm}^{-2} \text{ asec}^{-2}$) did target bright quasars. Bremer et al. (1992) finds extended emission around two out of three $z > 3$ quasars, but a search of another twelve quasars at similar redshifts revealed emission only around one source (Hu & Cowie 1987). Yet another search around a $z = 4.7$ quasar (Hu et al. 1996) has uncovered discrete Ly α emitting companions, rather than an extended continuous “fuzz”.

In summary, the few existing observations, typically utilizing ~ 1 hour integrations on 4m telescopes, probe surface brightnesses above $\gtrsim 10^{-16} \text{ erg s}^{-1} \text{ cm}^{-2} \text{ asec}^{-2}$, and reveal that luminous quasars are not generally enveloped by Ly α fuzz at this sensitivity. This puts a mild constraint on the models, although emission at this level is expected only in the largest halos (see Fig. 2). However, observations that probe bright quasars at about an order of magnitude deeper, would either detect Ly α fuzz around most sources, or else lead to strong constraints on the type of models discussed here. Provided that the Ly α fuzz is detected, its surface brightness, shape and extent, together with the line-profiles, would constitute an invaluable direct probe of galaxy formation.

It is interesting to consider constraints that would arise on galaxy and quasar formation, should the current trend of *not* detecting fuzz persist at fainter fluxes. One possibility is that quasars turn on only during the later stages of galaxy formation, i.e. at a time when most of the cold gas has already settled to a thin disk, and/or turned into stars. A lack of Ly α fuzz would then be naturally ex-

²See <http://www.subarutelescope.org>

plained by the absence of significant amounts of spatially extended cold gas during the luminous quasar phase. To avoid detectability at the surface brightness threshold of $\sim 10^{-17} \text{ erg s}^{-1} \text{ cm}^{-2} \text{ asec}^{-2}$, Figure 2 suggests that in halos with $T_{\text{vir}} \gtrsim 10^6 \text{ K}$, the flux has to be reduced by a factor $\gtrsim 10$ relative to the predictions of our simple models. This, implies, in turn, that $\gtrsim 90\%$ of the cold gas must already have settled to a disk (or disappeared). It has been argued that the presence of a disk is indeed a pre-requisite for the central BH to grow (Sellwood & Moore 1999); furthermore, cold gas in a thin disk is locally Toomre-unstable, and might rapidly turn into stars (e.g. Mo et al. 1998). Independent “evidence” favoring this scenario is that the heavy element abundances in the broad line region of quasars always appear to be high (Hamann 1999), even at high redshifts. This requires one or two generations of massive stars to precede the activation of the central quasar.

There are other, less attractive scenarios without Ly α emission. If the cold gas is quickly turned into stars, and/or collapse to the central regions, or to a thin disk, this could eliminate Ly α reprocessing, at least until cold gas is replenished by further accretion or merger with another halo (i.e. resulting in a short Ly α duty-cycle). However, this would also imply that a fraction $\sim t_{\text{dyn}}/t_Q \gtrsim 10\%$ of all quasars should still show Ly α fuzz, unless the cold gas disappears on an exceedingly short timescale ($\ll t_{\text{dyn}}$) (see Haiman & Hui for constraints on the lifetime t_Q of the luminous quasar phase). Alternatively, one might envision that the cold gas resides in clumps with a small covering factor, allowing most of the ionizing photons to leak out along line of sights traversing only hot (collisionally ionized) medium. However, this explanation requires a minimum cold clump size that exceeds the Jeans mass in the cold phase (Rees 1988). Hence, the postulated large clumps are unstable and would fragment to smaller pieces, increasing the covering factor to approximately unity. Yet another scenario with no Ly α fuzz is if the quasar (or its associated wind) has blown out most of the gas from the halo. However, in this case, one would still expect to see a fluorescent Ly α outflow around a fraction of quasars “caught in the act” of removing the gas.

Dust absorption might strongly suppress the Ly α flux escaping from a medium, even if the medium is optically thin in the Lyman continuum, and has been thought to cause the lack of detections of proto-galaxies in early Ly α surveys. A strong quenching of Ly α by dust, however, does not necessarily occur (see, e.g. Neufeld 1991, or Pritchett 1994 for a general discussion), especially if most of the Ly α photons originate in an ionized layer with relatively low dust opacity. Furthermore, Ly α emitting galaxies have been found at high-redshift (e.g. Hu et al. 1996; 1998), as expected in models with lower galactic dust abundance, and inhomogeneous dust distribution (Haiman & Spaans 1999). Based on these observations, it would appear unlikely that dust can suppress the Ly α fuzz from around all high-redshift quasars. Indeed, the dust abundance in the early, spatially extended, collapsing phase of the high-redshift halos is likely to be significantly lower than in star-forming galaxies.

Even if dust does not suppress the Ly α emission itself, however, an important question is whether the nucleus can

be obscured by dust and rendered undetectable at optical wavelengths. If this is typical, then Ly α fuzz should be expected around sub-mm sources, rather than around optical quasars. The extended Ly α emitting blob of Steidel et al. (2000) has been found to be an exceptionally bright sub-mm source (Chapman et al. 2000), with no visible continuum source. The Ly α line luminosity for this object is $\sim 2 \times 10^{10} L_{\odot}$, while the bolometric luminosity inferred from the sub-mm detections is $> 10^{13} L_{\odot}$. This implies that less than 1% of the UV produced by the source is available to power the extended Ly α emission, and that most of this UV is used up for this purpose (to explain the lack of any continuum source). It would be surprising if extended Ly α fuzz was typically produced in a similar way, since it requires a fine-tuning of the unobscured fraction to match the amount of surrounding cold gas. Nevertheless, it would be invaluable to target bright sub-mm sources (i.e. those with redshift estimates) in deep Ly α searches, to clarify what fraction of them do produce Ly α emission. The primary driver (stellar vs. quasar UV light) of both the sub-mm and Ly α emissions remain unclear in the Chapman et al. (2000) source (as well as in other bright sub-mm sources). Nevertheless, the lack of strong sub-mm emission in a second Ly α blob in the same dataset suggests that the line emission is not necessarily powered by a dust-obscured source.

Our models imply a second interesting, although somewhat less stringent constraint on the amount of extended cold gas around quasars. As discussed in § 3.1 above, we find that it is sufficient to convert $\sim 1\%$ of the total gas mass into a central BH in order to keep most of the cold gas ionized (cf. Eqn. 3). We emphasize that this must indeed happen if any ionizing radiation is to escape from the halo. Observations typically indicate that a large fraction of the ionizing radiation from quasars does escape, even for the quasar with the highest known redshift at $z = 5.8$ (Fan et al. 2000b). The abundance of this object implies a halo mass $M_{\text{halo}} \sim 10^{13} M_{\odot}$, while its luminosity, under the assumption that it equals the Eddington limit, implies a BH mass of $M_{\text{bh}} \sim 4 \times 10^9 M_{\odot}$ (Haiman & Loeb 2000). For this halo, under the assumption of a constant cold fraction $f = 0.5$, equation (3) would imply that a BH as large as $M_{\text{bh}} \sim 4 \times 10^{10} M_{\odot}$ is needed to ionize the cold gas and allow the ionizing radiation to escape. Based on the profile $f(r)$ derived in our models, the requisite BH mass in smaller, $M_{\text{bh}} \sim 1.3 \times 10^{10} M_{\odot}$. Nevertheless, this mass is close to 1% of the total gas mass, and is a factor of $\sim$ two higher than the BH mass inferred directly from the luminosity. We conclude that a large escape fraction of the ionizing continuum from bright, high-redshift quasars requires either (1) massive central BH’s whose masses are a significant fraction ($\sim 1\%$) of the gas mass, or (2) that some of the cold gas has settled to a disk (or disappeared in a blow-out).

Finally, we note that any significant continued gas in-fall at large radius, or a major merger spreading cold gas over an extended region, could still reprocess much of the quasar’s radiation to Ly α ; although the surface brightness at large radii might drop below detectable levels.

5. CONCLUSIONS

In this paper, we have studied what would happen to a protogalaxy if a bright quasar were to turn on dur-

ing the early stages of its assembly. Using a simple spherical model for the distribution of a two-phase gas, we find that such a system should have a substantial amount of cold, photoionized gas with a spatially extended distribution. We expect this conclusion to be generic, due to efficient radiative cooling at high redshifts. The cold, photoionized gas is detectable as an extended Ly α fuzz enshrouding the quasar; with a characteristic angular size of a few arcseconds, and a surface brightness of $\sim 10^{-18} - 10^{-16}$ erg s $^{-1}$ cm $^{-2}$ asec $^{-2}$. Existing observations have not yet targeted luminous quasars at these sensitivities, although at $\sim 10^{-16}$ erg s $^{-1}$ cm $^{-2}$ asec $^{-2}$, most quasars do not appear to be enveloped by extended emission. Future, deep Ly α imaging of few arcsecond regions around luminous quasars (as well as bright sub-mm sources) should be possible with current instruments.

While a detection of Ly α fuzz would provide a direct probe of galaxy formation; non-detections at the level of 10^{-17} erg s $^{-1}$ cm $^{-2}$ asec $^{-2}$ would already imply strong constraints.

We thank Eliot Quataert for useful discussions, and an anonymous referee for comments that improved this paper. This work began when MJR was visiting Princeton University as Scribner Lecturer; he acknowledges support from this source, and from the Royal Society. ZH was supported by NASA through the Hubble Fellowship grant HF-01119.01-99A, awarded by the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., for NASA under contract NAS 5-26555.

REFERENCES

- Baker, A. C., & Clements, D. L. 1997, in *Extragalactic Astronomy in the Infrared*, Eds. G. A. Mamon, T. X. Thuan, and J. Tran Thanh Van, Paris: Editions Frontieres, p. 295
- Bremer, M. N., Fabian, A. C., Sargent, W. L., Steidel, C. C., Boksenberg, A., & Johnstone, R. M. 1992, MNRAS, 258, 23p
- Böhringer, H., & Hensler, G., 1989, A&A, 215, 147
- Bunker, A. J., Marleau, F. R., & Graham, J. R. 1998, ApJ, 116, 2086
- Cavaliere, A., & Vittorini, V. 2000, ApJ, 543, 599
- Cen, R., & Haiman, Z. 2000, ApJL, 542, 75
- Chapman, S. C., Lewis, G. F., Scott, D., Richards, E., Borys, C., Steidel, C. C., Adelberger, K. L., & Shapley, A. E. 2000, ApJ, in press, astro-ph/0010101
- Cowie, L. L., & Hu, E. M. 1998, AJ, 115, 1319
- De Breuck, C., Röttgering, H., Miley, G., van Breugel, W., & Best, P. 2000, A&A, 362, 519
- Fabian, A. C. 1994, ARA&A, 32, 277
- Fall, S. M., & Efstathiou, G. 1980, MNRAS, 193, 189
- Fall, S. M., & Rees, M. J. 1985, 298, 18
- Fan, X., et al. 2000a, AJ, 119, 1
- Fan, X., et al. 2000b, AJ, 120, 1167
- Fardal, M. A., et al. 2000, ApJ, submitted, astro-ph/0007205
- Ferrarese, L., & Merritt, D. 2000, ApJL, 539, 9
- Gebhardt, K. et al. 2000, ApJL, 539, 13
- Gould, A., & Weinberg, D. 1996, ApJ, 468, 462
- Haiman, Z., & Hui, L. 2000, ApJ, 547, 27
- Haiman, Z., & Loeb, A. 2000, ApJL, in press, astro-ph/0011529
- Haiman, Z., & Spaans, M. 1999, ApJ, 518, 138
- Haiman, Z., Spaans, M., & Quataert, E. 2000, ApJL, 537, 5
- Hamann, F. 1999, ARA&A, 37, 487
- Heckman, T. M., Lehnert, M. D., Miley, G. K., & van Breugel, W. 1991a, ApJ, 370, 78
- Heckman, T. M., Lehnert, M. D., Miley, G. K., & van Breugel, W. 1991b, ApJ, 381, 373
- Hu, E. M., & Cowie, L. L. 1987, ApJ, 317, L7
- Hu, E. M., Cowie, L. L., & McMahon, R. G. 1998, ApJL, 502, 99
- Hu, E. M., McMahon, R. G., & Egami, E. 1996, ApJL, 459, 53
- Hu, E. M., Songaila, A., Cowie, L. L., & Stockton, A. 1991, ApJ, 368, 28
- Katz, N. 1991, ApJ, 368, 325
- Kauffmann, G., & Haehnelt, M. 2000, MNRAS, 311, 576
- Lacey, C., & Cole, S. 1993, MNRAS, 262, 627
- Madau, P., & Pozzetti, L. 2000, MNRAS, 312, 9
- Magorrian, J., et al. 1998, AJ, 115, 2285
- Makino, N., Sasaki, S., & Suto, Y. 1998, ApJ, 495, 555
- Menou, K., Haiman, Z., & Narayanan, V. 2000, ApJ, submitted
- Mo, H. J., Mao, S. & White, S. D. M. 1998, MNRAS, 295, 319
- Mo, H. J., & Miralda-Escudé, J. 1996, ApJ, 469, 589
- Moore, B., Ghigna, S., Governato, F., Lake, G., Quinn, T., Stadel, J. & Tozzi, P. 1999, ApJL, 524, 19
- Navarro, J. F., Frenk, C. S., & White, S. D. M. 1997, ApJ, 490, 493 [NFW]
- Navarro, J. F., & Steinmetz, M. 1997, ApJ, 478, 13
- Navarro, J. F., & White, S. D. M. 1994, MNRAS, 267, 401
- Neufeld, D. A. 1991, ApJ, 370, L85
- Osterbrock, D. E., 1989, *Astrophysics of Gaseous Nebulae and Active Galactic Nuclei*, University Science Books, Sausalito, CA
- Pritchett, C. J. 1994, PASP, 106, 1052
- Rees, M. J. 1984, ARA&A, 22, 471
- Rees, M. J. 1988, MNRAS, 231, 91
- Schneider, R., Ferrara, A., Ciardi, B., Ferrari, V., Matarrese, S. 2000, MNRAS, 317, 385
- Sellwood, J. A., & Moore, E. M. 1999, ApJ, 511, 660
- Steidel, C. C., Adelberger, K. L., Giavalisco, M., Dickinson, M., & Pettini, M. 1999, ApJ, 519, 1
- Steidel, C. C., Adelberger, K. L., Shapley, A. E., Pettini, M., Giavalisco, & M., Dickinson, M. 2000, ApJ, 532, 170
- van den Bosch, F. C., & Dalcanton, J. J. 2000, ApJ, 534, 146
- White, S. D. M., & Rees, M. J. 1978, MNRAS, 183, 341